

# A NONMEASURABLE SET FROM COIN FLIPS

ALEXANDER E. HOLROYD AND TERRY SOO

To motivate the elaborate machinery of measure theory, it is desirable to show that in some natural space  $\Omega$  one cannot define a measure on *all* subsets of  $\Omega$ , if the measure is to satisfy certain natural properties. The usual example is given by the Vitali set, obtained by choosing one representative from each equivalence class of  $\mathbb{R}$  induced by the relation  $x \sim y$  if and only if  $x - y \in \mathbb{Q}$ . The resulting set is not measurable with respect to any translation-invariant measure on  $\mathbb{R}$  that gives non-zero, finite measure to the unit interval [8]. In particular, the resulting set is not Lebesgue measurable. The construction above uses the axiom of choice. Indeed, the Solovay theorem [7] states that in the absence of the axiom of choice, there is a model of Zermelo-Frankel set theory where all the subsets of  $\mathbb{R}$  are Lebesgue measurable.

In this note we give a variant proof of the existence of a nonmeasurable set (in a slightly different space). We will use the axiom of choice in the guise of the well-ordering principle (see the later discussion for more information). Other examples of nonmeasurable sets may be found for example in [1] and [5, Ch. 5].

We will produce a nonmeasurable set in the space  $\Omega := \{0, 1\}^{\mathbb{Z}}$ . Translation-invariance plays a key role in the Vitali proof. Here shift-invariance will play a similar role. The **shift**  $T : \mathbb{Z} \rightarrow \mathbb{Z}$  on integers is defined via  $Tx := x + 1$ , and the shift  $\tau : \Omega \rightarrow \Omega$  on elements  $\omega \in \Omega$  is defined via  $(\tau\omega)(x) := \omega(x - 1)$ . We write  $\tau A := \{\tau\omega : \omega \in A\}$  for  $A \subseteq \Omega$ .

**Theorem 1.** *Let  $\mathcal{F}$  be a  $\sigma$ -algebra on  $\Omega$  that contains all singletons and is closed under the shift (that is,  $A \in \mathcal{F}$  implies  $\tau A \in \mathcal{F}$ ). If there exists a measure  $\mu$  on  $\mathcal{F}$  that is shift-invariant (that is,  $\mu = \mu \circ \tau$ ) and satisfies  $\mu(\Omega) \in (0, \infty)$ , and  $\mu(\{\omega\}) = 0$  for all  $\omega \in \Omega$ , then  $\mathcal{F}$  does not contain all subsets of  $\Omega$ .*

The conditions on  $\mathcal{F}$  and  $\mu$  in Theorem 1 are indeed satisfied by measures that arise naturally. A central example is the probability space  $(\Omega, \mathcal{G}, \mathbb{P})$  for a sequence of independent fair coin flips indexed by  $\mathbb{Z}$ , which is defined as follows. Let  $\mathcal{A}$  be the algebra of all sets of the form  $\{\omega \in \Omega : \omega(k) = a_k, \text{ for all } k \in K\}$ , where  $K \subset \mathbb{Z}$  is any finite subset of the integers and  $a \in \{0, 1\}^K$  is any finite binary string. The measure

$\mathbb{P}$  restricted to  $\mathcal{A}$  is given by  $\mathbb{P}(\{\omega \in \Omega : \omega(k) = a_k, \text{ for all } k \in K\}) = 2^{-|K|}$ , where  $|K|$  denotes the cardinality of  $K$ . Thus  $\mathbb{P}(\Omega) = 1$ , and  $\mathbb{P} = \mathbb{P} \circ \tau$  on  $\mathcal{A}$ . The Carathéodory extension theorem [6, Ch. 12, Theorem 8] gives a unique extension  $\mathbb{P}$  to  $\mathcal{G} := \sigma(\mathcal{A})$  (the  $\sigma$ -algebra generated by  $\mathcal{A}$ ) satisfying  $\mathbb{P} = \mathbb{P} \circ \tau$ . In addition, the continuity of measure implies  $\mathbb{P}(\{\omega\}) = 0$  for all  $\omega \in \Omega$ . Hence Theorem 1 implies that  $\mathcal{G}$  does not contain all subsets of  $\Omega$ . Of course, the same holds for any extension  $(\Omega, \mathcal{G}', \mathbb{P}')$  of  $(\Omega, \mathcal{G}, \mathbb{P})$  for which  $\mathbb{P}'$  is shift-invariant (such as the completion under  $\mathbb{P}$ ).

To prove Theorem 1 we will define a nonmeasurable function. We are interested in functions from  $\Omega$  to  $\mathbb{Z}$  that are defined everywhere except on some set of measure zero. Therefore, for convenience, introduce an additional element  $\Delta \notin \mathbb{Z}$ . Consider a function  $X : \Omega \rightarrow \mathbb{Z} \cup \{\Delta\}$ . We call  $X$  **almost-everywhere defined** if  $X^{-1}\{\Delta\}$  is countable, which implies that  $\mu(X^{-1}\{\Delta\}) = 0$ , for any measure  $\mu$  satisfying the conditions of Theorem 1. A function  $X$  is **measurable** with respect to  $\mathcal{F}$  if  $X^{-1}\{x\} \in \mathcal{F}$  for all  $x \in \mathbb{Z}$ . We call  $X$  **shift-equivariant** if

$$X(\tau\omega) = T(X(\omega)) \quad \text{for all } \omega \in \Omega$$

(where  $T(\Delta) := \Delta$ ). (We may think of a shift-equivariant  $X$  as an “origin-independent” rule for choosing an element from the sequence  $\omega$ .) Shift-equivariant functions of random processes are important in many settings, including percolation theory (for example in [2]) and coding theory (for example in [3, 4]).

**Lemma 2.** *If  $X : \Omega \rightarrow \mathbb{Z} \cup \{\Delta\}$  is an almost-everywhere defined, shift-equivariant function then  $X$  is not measurable with respect to any  $\mathcal{F}$  satisfying the conditions of Theorem 1.*

**Lemma 3.** *There exists an almost-everywhere defined, shift-equivariant function  $X : \Omega \rightarrow \mathbb{Z} \cup \{\Delta\}$ .*

Theorem 1 is an immediate consequence of the preceding two facts.

*Proof of Theorem 1.* Let  $(\Omega, \mathcal{F}, \mu)$  be a measure space satisfying the conditions of Theorem 1. Using Lemma 3, let  $X$  be an almost-everywhere defined shift-equivariant function. By Lemma 2,  $X$  is not  $\mathcal{F}$ -measurable. Therefore there exists  $z \in \mathbb{Z}$  such that  $X^{-1}\{z\} \notin \mathcal{F}$ .  $\square$

*Proof of Lemma 2.* Towards a contradiction, let  $X$  be a measurable function on  $(\Omega, \mathcal{F}, \mu)$  satisfying the conditions of Lemma 2. Since  $X$  is shift-equivariant we have for each  $x \in \mathbb{Z}$ ,

$$\mu(X^{-1}\{x\}) = \mu(\tau^{-x}X^{-1}\{x\}) = \mu(X^{-1}\{0\}).$$

Hence

$$\mu(X^{-1}\mathbb{Z}) = \mu\left(\bigcup_{x \in \mathbb{Z}} X^{-1}\{x\}\right) = \sum_{x \in \mathbb{Z}} \mu(X^{-1}\{0\}) = 0 \text{ or } \infty,$$

which contradicts the facts that  $\mu(X^{-1}\{\Delta\}) = 0$  and  $\mu(\Omega) \in (0, \infty)$ .  $\square$

Let us recall some facts about well-ordering. A total order  $\preceq$  on a set  $W$  is a **well order** if every nonempty subset of  $W$  has a least element. The well-ordering principle states that every set has a well order. It is a classical result of Zermelo [9] that the well-ordering principle is equivalent to the axiom of choice.

*Proof of Lemma 3.* Say  $\omega \in \Omega$  is **periodic** if  $\tau^x \omega = \omega$  for some  $x \in \mathbb{Z} \setminus \{0\}$ . If  $\omega$  is not periodic then  $(\tau^x \omega)_{x \in \mathbb{Z}}$  are all distinct. Using the well-ordering principle, fix a well order  $\preceq$  of  $\Omega$  and define the function

$$X(\omega) := \begin{cases} \Delta & \text{if } \omega \text{ is periodic;} \\ \text{the unique } x \text{ minimizing } \tau^{-x} \omega \text{ under } \preceq & \text{otherwise.} \end{cases}$$

(We may think of  $\tau^{-x} \omega$  as  $\omega$  viewed from location  $x$ , in which case  $X$  is the location from which  $\omega$  appears least.) Clearly,  $X$  is shift-equivariant. It is almost-everywhere defined since  $\Omega$  contains only countably many periodic elements.  $\square$

Acknowledgments. Alexander E. Holroyd is funded in part by an NSERC (Canada) Discovery Grant. Terry Soo is funded in part by an NSERC PGS D and a UBC Graduate fellowship.

## REFERENCES

- [1] D. Blackwell and P. Diaconis, A non-measurable tail set, in *Statistics, Probability and Game Theory*, IMS Lecture Notes-Monograph Series, vol. 30, Institute of Mathematical Statistics, Hayward, CA, 1996, 1–5.
- [2] R. M. Burton and M. Keane, Density and uniqueness in percolation, *Comm. Math. Phys.* **121** (1989) 501–505.
- [3] M. Keane and M. Smorodinsky, A class of finitary codes, *Israel J. Math.* **26** (1977) 352–371.
- [4] ———, Bernoulli schemes of the same entropy are finitarily isomorphic, *Ann. of Math. (2)* **109** (1979) 397–406.
- [5] J. C. Oxtoby, *Measure and Category*, 2nd ed., Graduate Texts in Mathematics, vol. 2, Springer-Verlag, New York, 1980.
- [6] H. L. Royden, *Real Analysis*, 3rd ed., Macmillan, New York, 1988.
- [7] R. M. Solovay, A model of set-theory in which every set of reals is Lebesgue measurable, *Ann. of Math. (2)* **92** (1970) 1–56.
- [8] G. Vitali, Sul problema della misura dei gruppi di punti di una retta, Gamberini and Parmeggiani, Bologna, 1905.

- [9] E. Zermelo, Beweis, daß jede Menge wohlgeordnet werden kann, *Math. Ann.* **59** (1904) 514–516.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, 121 –  
1984 MATHEMATICS RD, VANCOUVER, BC V6T 1Z2, CANADA.

*E-mail address:* holroyd@math.ubc.ca; tsoo@math.ubc.ca